



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

April 21, 2020

David Swain  
Authorized Agent for Activon, Inc.  
Scientific & Regulatory Consultants, Inc.  
201 W. Van Buren Street  
Columbia City, Indiana 46725

Subject: Label Amendment: Emerging Viral Pathogens Claim  
Product Name: EFFERSAN  
EPA Registration Number: 66570-2  
Application Date: March 19, 2020  
Decision Number: 561523

Dear Mr. Swain:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Because you have opted to add statements pertaining to emerging viral pathogens to your label as described in the August 19, 2016, Guidance to Registrants: Process For Making Claims Against Emerging Viral Pathogens Not On EPA-Registered Disinfectant Labels ("Guidance"), [https://www.epa.gov/sites/production/files/2016-09/documents/emerging\\_viral\\_pathogen\\_program\\_guidance\\_final\\_8\\_19\\_16\\_001\\_0.pdf](https://www.epa.gov/sites/production/files/2016-09/documents/emerging_viral_pathogen_program_guidance_final_8_19_16_001_0.pdf), you are subject to the following additional terms of registration:

1. You may make statements pertaining to emerging viral pathogens only through the following communications outlets: technical literature distributed exclusively to health care facilities, physicians, nurses and public health officials, "1-800" consumer information services, social media sites and company websites (non-label related). These statements shall not appear on marketed (final print) product labels.

2. Your statements pertaining to emerging viral pathogens must adhere to the format approved on the Agency-accepted master label.
3. You may make statements pertaining to emerging viral pathogens only upon a disease outbreak that meets all the following criteria:
  - a. The causative organism must be a virus that causes an infectious disease that has appeared in a human or animal population in the U.S. for the first time, or that may have existed previously but is rapidly increasing in incidence or geographic range.
    - i. For human disease, the outbreak is listed in one of the following Centers for Disease Control (CDC) publications:
      - A. CDC Current Outbreak List for “U.S. Based Outbreaks” ([www.cdc.gov/outbreaks](http://www.cdc.gov/outbreaks)),
      - B. CDC Current Outbreak List for “Outbreaks Affecting International Travelers” with an “Alert” or “Advisory” classification ([www.cdc.gov/outbreaks](http://www.cdc.gov/outbreaks)) (also released through the CDC’s Health Alert Network (HAN) notification process)
      - C. Healthcare-Associated Infections (HAIs) Outbreaks and Patient Notifications page ([www.cdc.gov/hai/outbreaks](http://www.cdc.gov/hai/outbreaks))
    - ii. For animal disease, the outbreak is identified as an infectious disease outbreak in animals within the U.S. on the World Organization for Animal Health (OIE) Weekly Disease Information page ([www.oie.int/wahis\\_2/public/wahid.php/Diseaseinformation/WI](http://www.oie.int/wahis_2/public/wahid.php/Diseaseinformation/WI)).
      - A. The CDC or OIE has identified the taxonomy, including the viral family and/or species, of the pathogen and provides notice to the public of the identity of the emerging virus that is responsible for an infectious disease outbreak. Based on the taxonomy of the outbreak pathogen identified by the CDC or OEI, the pathogen's viral subgroup are small non-enveloped, large non-enveloped, and enveloped.
      - B. The virus can be transmitted via environmental surfaces (non-vector transmission), and environmental surface disinfection has been recommended by the CDC, OIE or EPA to control the spread of the pathogen.
4. You may begin communicating statements pertaining to emerging viral pathogens only upon CDC or OIE’s publication per term 3.a. of an outbreak of an emerging viral pathogen meeting all of the criteria of term 3. You must cease and remove all such non-label communications intended for consumers no later than 24 months after the original publication of the outbreak per term 3.a., unless the Agency issue written guidance to the contrary due to continued public health concerns. The emerging pathogen claim language may remain on the master label.

5. Terms from points 1 through 4 above shall become immediately void and ineffective if registration for use against Feline Calicivirus and Canine parvovirus is suspended or cancelled or no longer meets the criteria for a disinfectant claim (see EPA Product Performance Test Guideline 810.2200). In addition, terms B.1 through B.4 above shall become immediately void and ineffective upon your receipt of evidence of ineffectiveness against any pathogen in a less-resistant Spaulding category.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, you may contact the disinfectants list at [disinfectantslist@epa.gov](mailto:disinfectantslist@epa.gov).

Sincerely,




Steven Snyderman, Acting Product Manager 33  
Regulatory Management Branch 1  
Antimicrobials Division (7510P)  
Office of Pesticide Programs

Enclosure: stamped label

## EfferSan™

Disinfectant, Bactericidal, Virucidal\*, Fungicidal, Sanitizer, Cleanser  
Protects Against Odor  
Eliminates and [ & ] Controls Odors  
Controls Bacteria and Algae  
Multi-purpose effervescent tablets  
Sanitizes, disinfects and protects against odor  
Sanitizes [kills] 99.999% [of] bacteria in 60 seconds  
The EfferSan Advantage  
Convenient accurate measurement  
EPA Registered  
Mild, non-irritating solution  
Safe [Appropriate] for Poultry, Cattle, Equine, Sheep Swine, Livestock Drinking Water  
[Consumption]  
Human Drinking Water Disinfectant



For use in Sanitation, Cleaning and Disinfection in amusement parks, breweries, beverage and food processing [plants], schools, hospitals, nursing homes, hotels, child care centers, daycares, restaurants, spas, hot tubs, salons, veterinary clinics, zoos and aquariums, milk processing facilities, dairy farms, farms, poultry premises [housing], poultry hatcheries, and livestock quarters, office buildings, industrial facilities, homes, camp sites, marine and recreational vehicles, R/V holding tanks, kennels, boarding facilities, laboratories, lab animal facilities, licensed care facilities.

Disinfects [precleaned, hard, nonporous surfaces] in 5 minutes.  
Effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella enterica* (formerly choleraesuis), \*Norovirus [(Feline Calicivirus as surrogate virus for Norovirus)]  
\*Canine parvovirus, \*Respiratory syncytial virus [(RSV)], \*2009-H1N1 Influenza A virus [(Novel H1N1)] [in 5 minutes]

Sanitizes precleaned, hard, non-porous food contact surfaces in 1 minute.  
No Rinse Sanitizer

At 200 ppm available chlorine, this product is an effective [food contact] [sanitizer] against *Campylobacter jejuni* [(ATCC 29428)], *Vibrio cholerae* (ATCC 11623), *Escherichia coli* O157:H7 [(ATCC 35150)], *Klebsiella pneumoniae* [(ATCC 4352)], *Listeria monocytogenes* [(ATCC 19117)], *Yersinia enterocolitica* [(ATCC 23715)], *Shigella sonnei* [(ATCC 25931)], *Salmonella typhi* (ATCC 6539) and *Staphylococcus aureus* [(ATCC 6538)] [in 1 minute]

At 1300 ppm available chlorine, this product is effective against *Trichophyton mentagrophytes* [(common cause of Athlete's Foot)] [in 5 minutes]

*{Front Panel continued}*

Active Ingredient:

Sodium Dichloro-s-Triazinetrione                      50%

Other Ingredients:    50%

Available Chlorine:    31.75%                      Total 100%

With *Activon* Technology

Keep Out of Reach of Children

**DANGER**

See back panel [inside] [booklet] for additional precautionary statements.

*{4g tablets}*

*{3.25g tablets}*

*{17.68g tablets}*

*{0.33g tablets}*

100 Tablets (0.88 lb.)    100 Tablets (0.72 lb.)    100 Tablets (3.9 lbs)    100 Tablets (0.073 lb.)

24 Tablets (0.21 lb.)

50 Tablets (1.95 lbs)

5 Tablets (0.044 lb.)

3 Tablets (0.423 oz)

*{Per PR Notice 98-10, additional package sizes and net contents may be added and are permitted by non-notification}*

EPA Reg. No. 66570-2

EPA Est. No. *{appropriate number to be inserted}*

[Manufactured for]

Activon, Inc.

123 Commercial Drive

Beaver Dam, WI 53916

*{Back Panel/Booklet}*

PRECAUTIONARY STATEMENTS  
 Hazards to Humans & Domestic Animals

DANGER: Corrosive: Causes irreversible eye damage. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing dust. Wear goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash separately before reuse.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment. [You may also call 1-800-222-1222 {or other appropriate number} for emergency medical treatment advice.]	
If in Eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If Swallowed	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor for treatment advice.</li> <li>• Have person drink large amounts of water if able to swallow. Avoid alcohol.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If Inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
If on Skin or Clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Immediately rinse skin with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.	

*{In accordance with PRN 2001-1, other formats may be used to present the first aid information}*

Physical or Chemical Hazards

Use only clean, dry utensils. Mix only into water. Contamination with moisture, dirt, organic matter or other chemicals (including pool chemicals) or any other foreign matter may start a chemical reaction with generation of heat, liberation of hazardous gasses and possible generation of fire and explosion. Avoid any contact with flaming or burning materials such as a lighted cigarette. Do not use this product in any chlorinating device that has been used with any inorganic or unstabilized chlorinating compounds (e.g., calcium hypochlorite). Such use may cause fire or explosion.

Note: Bracketed [ ] information is optional text. *{Bracketed Text}* is for administrative purposes only and will not appear on the printed label.

## Environmental Hazards

This pesticide is toxic to fish and aquatic organisms.

Disinfects floors, walls and other hard nonporous surfaces including walls, floors, tables, chairs, countertops, bathroom fixtures, sinks, shelves, racks, carts, refrigerators, coolers, glazed tile, linoleum, vinyl, glazed porcelain, plastic (such as polypropylene and polyethylene), stainless steel, or glass.

This product is designed for use in amusement parks, breweries, beverage and food processing plants, schools, hospitals, nursing homes, hotels, child care centers, restaurants, spas, hot tubs, veterinary clinics, zoos, milk processing facilities, dairy farms, farms, poultry premises, poultry hatcheries, and livestock quarters, office buildings, industrial facilities, homes, camp sites, marine and recreational vehicles, kennels, boarding facilities, laboratories, lab animal facilities.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## DISINFECTION

Prepare a 1,300 ppm solution [(refer to Dilution Chart)]. Clean surface then apply solution with mop, cloth, sponge, brush, [foaming equipment] or coarse trigger sprayer. Allow surface to remain wet for 5 minutes. Rinse thoroughly and allow to air dry. Prepare a fresh solution daily or when it becomes soiled or diluted. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

## SPA AND TUB USE

[Refer to Dilution Chart for number of tablets to use]

Add 4 ppm available chlorine [(refer to Dilution Chart)]. Using an appropriate test kit, test and adjust the water to the following values: pH: 7.2 – 7.8; total alkalinity: 60 – 100 ppm; calcium hardness: 200 ppm, minimum. Maintain these conditions for proper spa and hot tub operation by frequent testing with a test kit. Do not allow cyanuric acid level to exceed 150 ppm. It is recommended that spas and hot tubs be drained every 60 – 90 days, more often under heavy use. Consult manufacturer's recommendations concerning the compatibility of chlorine sanitizers with their equipment. Some oils, lotions, fragrances, cleansers, etc., may cause foaming or cloudy water and may react with chlorine sanitizers, reducing their efficacy. Reentry into treated spas/hot tubs is prohibited above levels of 3 ppm chlorine.

Note: Bracketed [ ] information is optional text. {Bracketed Text} is for administrative purposes only and will not appear on the printed label.

#### Start-Up (Freshly Filled)

1. Turn on the circulation system and ensure that it is operating properly.
2. Add 4 ppm available chlorine [(refer to Dilution Chart)]. Check the free available chlorine (FAC) level and, if below 4 - 5 ppm, repeat as needed.

#### Regular Use

Turn on the circulation system and ensure that it is operating properly. Add 4 ppm available chlorine [(refer to Dilution Chart)] to the water. Test for FAC and add additional product, if necessary, to attain 4-5 ppm FAC. Maintain 1 – 3 ppm FAC while the spa or hot tub is in use. After each use, shock treat with 10 ppm available chlorine [(refer to Dilution Chart)] to control odors and algae. Repeat as needed. Spa or hot tub should not be entered until FAC reaches 1 – 3 ppm.

#### Extended Non-Use Period

During extended periods when the spa or hot tub is not being used, with the circulation system running, add 4 ppm available chlorine [(refer to Dilution Chart)] twice a week or as needed to maintain 1 – 3 ppm FAC.

{FOR USE IN removed} INDUSTRIAL RECIRCULATING WATER COOLING TOWERS, AIR WASHERS & EVAPORATIVE CONDENSERS

[Refer to Dilution Chart for number of tablets to use]

Treatment with this product is an effective way to control the growth of bacteria and algae in industrial recirculating water cooling towers, air washers and evaporative condensers.

1. Clean badly fouled systems prior to initiating treatment.
2. Initial Dosage – when the system is just noticeably fouled, add 3 ppm available chlorine [(refer to Dilution Chart)] to the system water. Repeat this dosage, if necessary, until a free available chlorine (FAC) level of 0.5 – 1.0 ppm is obtained, as determined by use of a reliable test kit.
3. Maintenance Dosage – to obtain a FAC of 0.5 – 1.0 ppm, add 0.5 ppm -1.0 ppm available chlorine [(refer to Dilution Chart)] daily or as needed.
4. Add this product to the system at a point where adequate flow is maintained. Variations in water temperature, chlorine demand and flow rate will affect the dissolution rate. Warmer seasons may require an upward adjustment of the FAC.

{AIR WASHERS section removed}

{FOR USE IN removed} SEWAGE TREATMENT [Not applicable in California]

1. Disinfection of Effluents: Disinfection does not occur instantaneously. A suitable detention basin must be provided to expose the sewage effluent to the effects of this product for a sufficient period of time (usually a minimum of 15 minutes). Where mechanical stirring or other agitation is not present, introduce product solution before primary or secondary sedimentation treatments, if these are used.

The amount of product solution required will vary, depending on the concentration and conditions of the final effluent. Treat the sewage before it has reached a septic state. About

30% of the chlorine demand of raw sewage is attributed to settle solids; 40% to suspended and colloidal solids; and 30% to dissolved solids.

Whenever possible, control disinfection by laboratory checks. Disinfection can be achieved when the chlorine residual (after 15 – 30 minutes contact time) is between 0.6 and 1.0 ppm. Experience with different types of treated sewage may eventually establish a relationship between the residual chlorine content of the final effluent and the contact time necessary to ensure the desired bacteriological results. Once this relationship is established, the residual chlorine content and contact time may then become the controlling factors for operation. Perform occasional bacteriological checks as a safeguard.

In cases where sewage is to be temporarily disinfected before being diluted in a body of water, the following conditions will usually provide satisfactory protection against receiving waters' pollution:

- a. Raw sewage: 10 – 30 ppm available chlorine
- b. Primary treated sewage: 5 – 20 ppm available chlorine
- c. Sewage which has undergone primary and secondary treatment, or secondary alone: 2 – 5 ppm

Frequently perform bacteriological tests as a safeguard. The available chlorine level in the discharge effluent should be between 0.6 and 1.0 ppm or in accordance with an NPDES permit. For guidance contact the regional office of the EPA.

2. Slime Control: When ponding of the filters is excessive, stoppage of the distributing filter can occur. Add 10 ppm available chlorine [(refer to Dilution Chart)] into the effluent at a point above the filter nozzles. Repeat as necessary until the desired cleaning has been achieved. To maintain the system, intermittently apply a solution of this product to the dosing tanks, just ahead of the filter. The amount and frequency of the dosage needed to give satisfactory continuous operation of the trickling filters depends on the severity of the microbiological problem.
3. Biological Oxygen Demand (B.O.D.) Reduction: The condition can usually be avoided by applying the product solution to the effluent until a substantial residual is obtained. Apply at a point that will permit 10 – 20 minute contact time prior to discharging effluent into the stream. A dosage that leaves a residual available chlorine of about 0.2 ppm after a contact time of at least 10 minutes will afford a reduction of about 1/3 of the effluents B.O.D. Where more permanent or greater B.O.D. reduction is necessary dosing to high available chlorine residuals is recommended.
4. Coagulation and Sedimentation: A great deal of the finer divided suspended matter and most of the colloidal matter in sewage does not readily respond to plain sedimentation. The job of removing substantial portions of this kind of matter is usually accomplished either by chemical precipitation, by filtration or by the use of both processes. Chlorine improves sedimentation and coagulation in sewage treatment operations.

5. Treating Effluent from Mobile Sewage Treatment Units (Including Marine and Recreational Vehicles): Only human waste, toilet paper and water should enter the mobile sewage treatment unit. Solids are retained in the unit for later removal, while the liquid portion is filtered, disinfected and discharged. Product is placed in a flow-through container where the liquid effluent passes over them before being discharged.

Disinfection does not occur instantaneously. A suitable detention basin must be provided to expose the sewage effluent to the effects of this product for a sufficient period of time (usually a minimum of 15 minutes). Frequently test effluent as a safeguard. The available chlorine level in the discharge effluent should be between 0.6 and 1.0 ppm or in accordance with an NPDES permit. [Refer to Dilution Chart for number of tablets to use]

#### FOR USE THROUGHOUT FOOD AND BEVERAGE PROCESSING AND FOOD HANDLING OPERATIONS

[Refer to Dilution Chart for number of tablets to use]

This product is recommended for sanitizing all types of hard, non-porous equipment and utensils used in food processing and canning plants, bottling plants, breweries, fish processing plants, meat and poultry processing plants, milk handling and processing plants, restaurant and institutional dining establishments. Use a 100 ppm available chlorine solution [(refer to Dilution Chart)] to sanitize previously cleaned processing and packaging equipment. Allow at least a one minute contact time before draining. Allow adequate draining before contact with beverages.

To control the growth of bacteria in brewery pasteurizers, clean badly fouled systems before treatment. When the system is noticeably fouled, add 3 ppm available chlorine [(refer to Dilution Chart)] to system water. Repeat this dosage if necessary until the free available chlorine (FAC) level is 0.5 – 1.0 ppm, as determined by use of a reliable test kit. To maintain a FAC of 0.5 – 1.0 ppm, add 0.5 ppm – 1 ppm available chlorine [(refer to chart)] daily as needed. Add this product to the system at a point where adequate flow is maintained.

#### EGG PROCESSING PLANTS

[Refer to Dilution Chart for number of tablets to use]

Clean and destain egg shells prior to sanitizing. To clean egg shells, spray with a 90°F to 120°F 100 ppm available chlorine solution [(refer to Dilution Chart)]. Spray-rinse the cleaned eggs with warm potable water. Only clean, whole eggs may be sanitized. Dirty, cracked or punctured eggs may not be sanitized.

To destain egg shells, immerse the eggs in a 90°F to 120°F solution containing 100 ppm available chlorine [(refer to Dilution Chart)]. After destaining, the eggs must be cleaned by spraying with an acceptable cleaner. Follow with a potable water rinse.

To sanitize clean shell eggs intended for food or food products, spray with a solution containing 100 ppm available chlorine [(refer to Dilution Chart)]. The solution must be equal to or warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs that have been sanitized with this chlorine compound may be broken for use in the manufacture of egg products without a prior potable water rinse. Eggs must be reasonably dry before casing or breaking. The solution must **not** be reused for sanitizing eggs.

Note: Bracketed [ ] information is optional text. {Bracketed Text} is for administrative purposes only and will not appear on the printed label.

Thoroughly clean and sanitize all egg cups, breaking knives, trays and other equipment that come into contact with “off” eggs. First, clean all equipment. Before placing back in use, spray with a solution containing 100 ppm available chlorine [(refer to Dilution Chart)]. Allow surfaces to completely drain before contact with egg product. To sanitize egg freezers and dryers (tanks, pipelines and pumps), use the spray method of treatment (**see Sanitizing Application Methods section**). This procedure is generally used to sanitize large, non-porous surfaces that have already been cleaned of physical soil.

Prepare a solution containing 100 ppm available chlorine [(refer to Dilution Chart)]. Heavily apply spray to all surfaces the eggs will touch. Thoroughly spray all treated surfaces, corners and turns. Allow at least a one minute contact time before draining. Allow equipment to drain adequately before contact with eggs.

#### SANITIZING HARD, NON-POROUS SURFACES, DISHES, GLASSES, FOOD PROCESSING EQUIPMENT AND UTENSILS, DAIRY AND BREWERY EQUIPMENT AND UTENSILS [Refer to Dilution Chart for number of tablets to use]

This product is an effective sanitizing agent. Treatment with this product throughout food and beverage processing and food handling operations can help ensure the quality and safety of the final product.

#### Hand Washing of Items

1. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution.
2. Prepare a 100 ppm available chlorine sanitizing solution [(refer to Dilution Chart)].
3. Place equipment, utensils, dishes, glasses, etc. in the solution or apply the use solution to surfaces using a cloth, sponge, or coarse sprayer.
4. Allow to stand at least one minute, drain the excess solution from the surface and allow to air dry.
5. Fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

#### Machine Washing of Items

1. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution.
2. Prepare a 100 ppm available chlorine solution [(refer to Dilution Chart)].
3. Add the solution to the feed tank of immersion or spray type machines that can provide at least one minute contact time for sanitizing dishes, glasses, food processing equipment or utensils. Allow to drain and air dry before use.
4. Promptly use the sanitizing solution. Prepared solutions cannot be reused for sanitizing but may be used for other purposes, such as cleaning.

#### ANIMAL HOUSING FACILITIES (Including Poultry Houses, Swine Confinement Facilities, Veterinary Clinics, Zoos and Farms)

Note: Bracketed [ ] information is optional text. {Bracketed Text} is for administrative purposes only and will not appear on the printed label.

The problem of odor control in poultry houses and other animal facilities is not completely solved by normal cleaning practices. The regular use of an efficient bactericide and deodorant is strongly recommended and often required by health authorities.

Remove all poultry or animals and feeds from premises, trucks, vehicles, coops, crates and enclosures. Remove all litter and manure or droppings from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. To disinfect, saturate all surfaces with a 1,300 ppm available chlorine solution [(refer to Dilution Chart)] for a period of five minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, coops and other closed spaces. Do not house livestock or poultry or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

#### SHOE AND BOOT BATH SANITIZER [Not applicable in California]

[To prevent cross contamination into treated animal areas and the packaging and storage areas of food plants.] Shoe and Boot baths containing one inch of freshly made 100 ppm available chlorine [(refer to Dilution Chart)] should be placed at all entrances to buildings, hatcheries and at all the entrances to the production and packaging rooms. Scrape waterproof [shoes] [and] [boots] and place into solution for at least one minute [60 seconds] prior to entering area. Change the sanitizing solution in the bath at least daily or sooner if solution appears diluted or dirty.

#### MILK HANDLING AND PROCESSING EQUIPMENT

This product can be used on dairy farms and in plants processing milk, cream, ice cream and cheese. Rinse milking machines, utensils and all equipment with cold water to remove excess milk. Clean and rinse prior to sanitizing. To sanitize, spray or rinse all precleaned surfaces with 100 ppm available chlorine [(refer to Dilution Chart)]. Allow adequate draining before contact with dairy products.

It is important to clean out large deposits of milk or other organic matter before sanitizing. A sharp decline in the available chlorine content of the sanitizer following circulation through milk processing equipment is usually regarded as evidence of inadequate cleaning of the equipment and should be promptly investigated.

#### SANITIZING APPLICATION METHODS

Freshly prepare all sanitizing solutions. Test solutions during use to ensure the concentration does not drop below the recommended level. Keep in properly labeled containers to protect against contamination. Discard unused solutions.

#### **Pressure Method of Sanitizing Equipment**

Note: Bracketed [ ] information is optional text. {Bracketed Text} is for administrative purposes only and will not appear on the printed label.

This method is commonly used to sanitize closed systems, such as fluid milk cooling and handling equipment. It is also appropriate for sanitizing weigh tanks, coolers, short-time pasteurizers, pumps, homogenizers, fillers, sanitary piping and fittings, and bottle and can fillers. For mechanical operations, prepared solutions cannot be reused for sanitizing but may be used for other purposes, such as cleaning. For manual operations, fresh sanitizing solutions must be prepared at least daily or more often if the solution becomes diluted or soiled. First, disassemble and thoroughly clean all equipment immediately after use. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution. Then place back in operating position. Prepare a solution containing 100 ppm available chlorine [(refer to Dilution Chart)] in a volume equal to 110% of capacity. Pump the solution through the system until it is filled with sanitizer and air excluded. Close final drain valves and hold under pressure for one minute to ensure proper contact with all surfaces. Remove a portion of the cleaning solution from the drainvalve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine. {"Rinse system with potable water prior to use" removed.}

### **Spray Method of Sanitizing Equipment**

The spray method is generally used to sanitize large, non-porous surfaces that have already been freed of physical soil. It is appropriate for batch pasteurizers, holding tanks, weigh tanks, tank trucks and cars, vats, tile walls, ceilings and floors. Clean all surfaces after use. Prepare a solution containing 100 ppm available chlorine [(refer to Dilution Chart)]. Use pressure spraying equipment designed to resist chlorine-containing solutions (e.g. rubber-coated, plastic or stainless steel). When using any other kind of spraying equipment, always empty and thoroughly rinse the spray equipment with potable water immediately after treatment. Apply spray heavily to all surfaces the product will touch. Thoroughly spray all treated surfaces, corners and turns until wet. Allow at least a one minute contact time before draining. Allow excess solution to drain and air dry then place in service. Vacate area for at least two hours.

### **General Rinse Method**

Solutions containing 100 ppm available chlorine [(refer to Dilution Chart)] sanitize plant floors, walls and ceilings, and also control odors in refrigerated areas and drain platforms. Generously flush or swab surfaces with the solution. After one minute contact time allow solution to drain and then air dry.

## **DISINFECTION OF DRINKING WATER**

### **DRINKING WATER**

Add 10 ppm available chlorine [(refer to Dilution Chart)] to water to be disinfected. Let the water stand seven to fifteen minutes before using. A residual of 0.2 ppm available chlorine, as measured by a reliable test kit, should be maintained in the water to insure disinfection.

### **EMERGENCY DRINKING WATER**

This product may be used to disinfect or pre-treated (settled, coagulated and/or filtered) water supplies intended for use as drinking water for humans and domestic animals on an emergency basis as defined in 40 CFR, Part 165-179.

The source of the water to be treated may be a river, lake, well, cistern or similar system. To obtain the desired disinfection results, the water to be treated should be clear and free dirt and organic debris. If the source of the water is cloudy and contains dirt and organic debris, the water should be held in holding tanks or ponds, treated with coagulating agents and filtered to remove the dirt and organic debris.

Preparation of Stock Solution – Dissolve proper amount of this product to produce a 6,000 ppm available chlorine [(refer to Dilution Chart)] stock chlorine solution. Add 20 drops of this stock solution for each liter of water to be treated. The stock solution should be prepared fresh weekly.

## **EMERGENCY DISINFECTION AFTER FLOODS**

### **DISTRIBUTION SYSTEM**

Flush repaired or replaced section with water. Establish a chlorinating station and apply sufficient product until a consistent available chlorine residual of at least 10 ppm available chlorine (as measured by a chlorine test kit) remains after a 24 hour retention time. [(refer to Dilution Chart)]

## **EMERGENCY DISINFECTION AFTER DROUGHT**

### **WATER SHIPPED IN BY TANKS, TANK CARS, ETC.**

Thoroughly clean all containers and equipment. Spray a 500 ppm available chlorine solution [(refer to Dilution Chart)] and rinse with potable water after 5 minutes. During the filling of the containers, dose with sufficient amounts of this product to provide at least a 0.2 ppm chlorine residual, as measure by a chlorine test kit.

## **POULTRY, SWINE, CATTLE, LIVESTOCK DRINKING WATER DISINFECTION**

Effersan [This product] can be used as a disinfectant for the drinking water of poultry, swine, cattle and other livestock. If the water supply is badly fouled, then add 5 ppm available chlorine [(refer to Dilution Chart)] to the water supply. After 24 hours the addition rate can be reduced to 1 ppm available chlorine [(refer to Dilution Chart)]. If the microbiological content of the water is eliminated, the concentration of available chlorine can be reduced to 0.5 ppm. If the microbiological control is not adequate at 1 ppm available chlorine, then add 1.5 ppm available chlorine to the livestock drinking water.

Effersan [This product] should be administered continuously into the drinking water from the time of placement (day one). Remove Effersan [this product] from the drinking water 24 hours prior to vaccination and re-administer 24 hours after vaccination.

## **FILTRATION DEVICES**

This product is for use in filtration devices (water purification systems and its cartridges). Its purpose is to clean membranes, such as reverse osmosis membranes of fouling contaminants. [One four [4] gram tablet] [One [1] dose] of product is necessary to achieve cleaning of fouling contaminants. Product should be used following the manufacturer's instructions.

*{The following text is optional for "One dose" option above}*

[Add 3 ppm available chlorine [(refer to Dilution Chart)] to the system water. Repeat this dosage, if necessary, until a free available chlorine (FAC) level of 0.5 – 1.0 ppm is obtained, as determined by use of a reliable test kit.]

{the appropriate chart will be used on the market label for each tablet size and number of tablets in container}

Dilution Chart {4 g Tablets}			Dilution Chart {0.33g tablets}			Dilution Chart {17.68g tablets}			Dilution Chart {3.25g tablets}		
Solution (Available chlorine)	Tablets (4g or 0.14 oz. each)	Gallons of Water	Solution (Available chlorine)	Tablets (0.33g or 0.01oz each)	Gallons of Water	Solution (Available chlorine)	Tablets (17.68g or 0.62oz each)	Gallons of Water	Solution (Available chlorine)	Tablets (3.25g or 0.11 oz each)	Gallons of Water
0.5ppm	1	625	0.5ppm	1	55				0.5ppm	2	625
	4	2,500		10	550	0.5ppm	1	2,500		5	2,500
	8	5,000					2	5,000		10	5,000
	15	10,000					4	10,000		19	10,000
1ppm	1	325	1ppm	1	25				1ppm	2	325
	2	650		5	135					3	650
	3	1,000		10	275	1ppm	4	6,000		4	1,000
	30	10,000					7	10,000		37	10,000
3 ppm	1	100	3 ppm	11	100				3 ppm	2	100
	5	500		25	230	3 ppm	1	500		6	500
	9	1,000		55	500		2	1,000		11	1,000
	90	10,000					20	10,000			
4 ppm	6	500	4 ppm	73	500				4 ppm	8	500
	15	500		2	5	4 ppm	3	1,000		19	500
10 ppm	1	3	10 ppm	1	1 quart				10 ppm	2	3
	3	10		4	1	100 ppm	1	15		4	10
100 ppm	12	40							100 ppm	15	40
	4	1	1,300ppm	12	1 quart					5	1
1,300ppm	39	10		47	1				1,300ppm	48	10

Note: Bracketed [ ] information is optional text. {Bracketed Text} is for administrative purposes only and will not appear on the printed label.

## Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Keep product dry in tightly closed container when not in use. Store in a cool, dry, well-ventilated area away from heat or open flame.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

*{for tablets in plastic container}*

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Thoroughly rinse empty container with water to dissolve all material before discarding in trash or recycling.

*{for individually wrapped tablets in paper box}*


Container Disposal: Throw tablet wrapper in trash. Outer container – Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or dispose in trash.

*{for individually wrapped tablets in plastic container}*

Container Disposal: Throw tablet wrapper in trash. Outer container – Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Thoroughly rinse empty container before discarding in trash or recycling.

## Emergency Handling

In case of contamination or decomposition do not reseal container. If possible, isolate container in open and well-ventilated area. Flood with large volumes of water. Dispose of contaminated material in an approved landfill area.



{Additional Graphic above for 4 gram tablets}

**{Emerging Viral Pathogens}**

{This product qualifies for emerging viral pathogen claims per the EPA’s ‘Guidance to Registrants: Process for Making Claims Against Emerging Viral Pathogens not on EPA-Registered Disinfectant Labels’ when used in accordance with the appropriate use directions indicated below.

The statements shall be made only through the following communications outlets: technical literature distributed exclusively to health care facilities, physicians, nurses and public health officials, "1-800" consumer information services, social media sites and company websites (non-label related). These statements shall not appear on marketed (final print) product labels.}

<i>For an emerging viral pathogen that is a/an...</i>	<i>...follow the directions for use for the following organisms on the label:</i>
Enveloped virus	Norovirus (Feline Calicivirus as surrogate) [(ATCC VR-782)]
Large, non-enveloped virus	Norovirus (Feline Calicivirus as surrogate) [(ATCC VR-782)]
Small, non-enveloped virus	Norovirus (Feline Calicivirus as surrogate) (ATCC VR-782) & Canine parvovirus (ATCC VR-2017)

This product [-or- Product Name] has demonstrated effectiveness against viruses similar to {Insert name of emerging virus} on hard, nonporous surfaces. Therefore, this product [-or-Product Name] can be used against {Insert name of emerging virus} when used in accordance with the directions for use against Norovirus (Feline Calicivirus as surrogate) and Canine parvovirus on hard, nonporous surfaces. Refer to the CDC -or- OIE website at {Insert pathogen-specific website address} for additional information.

{Insert name of illness/outbreak} is caused by {Insert name of emerging virus}. This product [-or-Insert Name] kills similar viruses and therefore can be used against {Insert name of emerging virus} when used in accordance with the directions for use against Norovirus (Feline Calicivirus as surrogate) and Canine parvovirus on hard, nonporous surfaces. Refer to the {CDC -or- OIE} website at {Insert pathogen-specific website address} for additional information.

Note: Bracketed [ ] information is optional text. {Bracketed Text} is for administrative purposes only and will not appear on the printed label.